

Indirect chemiluminescent immunoassay (CLIA) to detect antibodies against SARS-CoV-2, producing COVID-19, in human serum or plasma.

COVID-19 VIRCLIA® IgM+IgA MONOTEST COVID-19 VIRCLIA® IgG MONOTEST

- · SARS-CoV-2 recombinant antigens.
- · Simple and automated protocol with results in 1 hour.
- · Unique and definitive solution for urgent samples.
- · Same day results, no batching, no sample cumulation.
- · Complete solution: IgG and IgM+IgA kits available.
- Simultaneous detection of IgM and IgA improves sensitivity in the acute phase of the disease.
- · Excellent sensitivity with proven PCR positive samples.



VirClia® MONOTEST - 24 tests



THE TRUE MONOTEST

SARS-CoV-2. Vircell culture collection

- On-demand testing: One monotest= one reportable test. Nothing else is required.
- SARS-CoV-2 recombinant antigens: spike glycoprotein (S) and nucleocapside protein (N).
- Individual Quality Control per monotest, no need for extra controls or calibrations.



Excellent performance in samples characterized by PCR

IgG

Sensitivity 92% in patients with more than 9 days after a positive PCR result

Specificity 99% in samples from healthy pre-pandemic donors

IgM+IgA

78% in patients 5 days after a positive PCR result

99% in samples from healthy pre-pandemic donors



Random Access, Continuous Loading.



Total flexibility, STAT function.



Primary sample tubes.
Positive identification of samples & strips.

Ordering information

Reference	Description	Pack size
VCM097	COVID-19 VIRCLIA® IgG MONOTEST	24 tests
VCM098	COVID-19 VIRCLIA® IgM+IgA MONOTEST	24 tests

Telefon: +49 2234 98795 - 0 Telefax: +49 2234 98795 - 29 Email: service@bestbion.com Internet: www.bestbion.com bestbion dx GmbH Horbeller Str.33 50858 Köln Deutschland

